



Test Update 836

Posted Date 03/15/2023
Effective Date 03/16/2023
Test Name [Cortisol, Saliva \(SCOR\)](#)
Update Type [New Tests](#)

NEW TEST

Cortisol, Salivary

Order Code: SCOR
Fee Code: KA055 (CPT 82533)

MLabs Toxicology Laboratory will begin offering Salivary Cortisol testing by liquid chromatography mass spectrometry (LC-MS/MS) effective February 28, 2023. This test replaces Mayo Clinic Laboratories test ID SALCT.

Measurement of midnight/late-night salivary cortisol (LNCS) is used for screening of adrenal hyperfunction (Cushing syndrome). Although other cortisol tests may be ordered for the initial workup, LNCS has a superior sensitivity to plasma and 24-hour urinary free cortisol measurements. A positive LNCS result (≥ 100 ng/dL) should be followed by a second test for cortisol which may include LNCS, 24-hour urinary cortisol or dexamethasone suppression testing. Two positive LNCS may be used to confirm Cushing Syndrome.

The order code SCOR will reflex to one of the following dependent on the time of collection:

Code	Collection Time	Reference Range	LOINC
ACOR	0700 - 0800	100-750 ng/dL	51494-3
PCOR	1500 - 1700	<401 ng/dL	51493-5
MCOR	2300 - 2400	<100 ng/dL	51492-7
SACOR	any time outside of the time frames above		51495-0

Collection Instructions: The preferred time of collection is between 2300 - 2400. Collect saliva specimen using Starstedt Salivette Cortisol Saliva Collection Device. Because of the onset of bacterial growth in saliva after a few hours at room temperature, it is recommended to either centrifuge the firmly closed Salivette® Cortisol within 4 hours after specimen collection at 1,000 x g for 2 minutes. Alternatively store and transport refrigerated (unspun).

Patient Collection Instructions:

The saliva sample should be taken at least 60 minutes after a meal (liquid/solid food intake), brushing teeth or taking oral medication, in order to avoid the saliva being contaminated with interfering substances. Up to 10 minutes before taking the sample, rinse mouth with water for 1-5 seconds. Immediately before collecting the saliva, wash and dry hands. With two fingers, remove the absorbent roll at the far end from the opened Salivette® Cortisol and place it in the mouth. The absorbent roll should be kept in the mouth (e.g., in the cheek) for 2 minutes without chewing. With two fingers, take the absorbent roll out of the mouth by the far end and place it immediately in the Salivette® Cortisol and seal with the blue cap. Store refrigerated.

[View PDF](#)