



## **Test Update 848**

**Posted Date** 07/05/2023

**Effective Date** 07/03/2023

**Test Name** [Matrix Metalloproteinase-7 \(MMP-7\)](#)

**Update Type** [Miscellaneous](#)

### **PERFORMING LABORATORY CHANGE**

#### **Matrix Metalloproteinase-7 (MMP-7)**

Order Code: CMMP7

Fee Code: AB050 (CPT 83519)

New Fee Code: pending (CPT 83520)

Reference Laboratory: Cincinnati Children's 11738598

New Reference Lab: Children's Medical Center of Dallas

Cincinnati Children's has discontinued the Matrix Metalloproteinase-7 (MMP-7) assay effective June 30, 2023. Requests for this test will be sent to Children's Medical Center, Dallas, TX. There will be no change to the order code.

Collection Instructions: Collect specimen in an SST or red top tube. Place on ice immediately after collection. Centrifuge, aliquot serum into a plastic vial and freeze within 2 hours of collection.

Reference Range: 3.6 - 10.5 ng/mL

MMP-7 level  $\geq 18.2$  ng/mL has 95% sensitivity and 65% specificity for diagnosis of biliary atresia, with negative predictive value of 97%.

Caution should be exercised in the interpretation of any assay for MMP-7, and the results should be considered in conjunction with the patient's clinical condition. High MMP-7 have been detected in premature babies and young infants with cardiovascular and/ or multi-system disease. The final diagnosis of biliary atresia requires intraoperative cholangiography and the histological presence of obstruction of the extrahepatic bile duct. The normal values of MMP-7 in premature babies have not yet been established. Clinical correlation is particularly important in premature babies.

[View PDF](#)

2800 Plymouth Rd, Bldg 35 • Ann Arbor, MI 48109-2800 • 800.862.7284 • 734.936.0755 Fax • [mlabs.umich.edu](mailto:mlabs.umich.edu)